

Rycar®

Insecticide

SPECIMEN



GROUP UN INSECTICIDE

For Indoor Greenhouse Use on Ornamental Plants
Not for residential use sites

Active Ingredient:

Pyrifluquinazon [1-acetyl-1,2,3,4-tetrahydro-3-[(3-pyridylmethyl)amino]-6-[1,2,2,2-tetrafluoro-1-(trifluoromethyl)ethyl]quinazolin-2-one].....20.0%

Other Ingredients80.0%

TOTAL.....100.0%

Contains 1.80 lbs. active ingredient per U.S. gallon

Keep Out of Reach of Children

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Refer to label booklet for additional Precautionary Information and Directions for Use, including Storage and Disposal.

Notice: Read the entire label before using. Use only according to label directions. Before buying or using this product, read *Terms and Conditions of Use, Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies* inside label booklet. If terms are unacceptable, return at once unopened.

®Rycar is a registered trademark of Nichino America, Inc. EPA Reg. No. 71711-37-67690
Manufactured for: **SePRO Corporation** 11550 North Meridian Street FPL20131216
Suite 600, Carmel, IN 46032 U.S.A.

FIRST AID

| | |
|-------------------------------|--|
| If on skin or clothing | <ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15-20 minutes.Call a poison control center or doctor for treatment advice. |
| If swallowed | <ul style="list-style-type: none">Call poison control center or doctor immediately for treatment advice.Have person sip a glass of water if able to swallow.Do not induce vomiting unless told to do so by the poison control center or doctor.Do not give anything to an unconscious person. |

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC at 1-800-535-5053**.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or the toilet. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves. Remove and wash contaminated clothing before reuse.

Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, made of material such as barrier laminate or Viton
- Shoes plus socks

For more options for chemical-resistant gloves, follow the instructions for Category C on an EPA chemical-resistance category selection chart.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

User Safety Recommendations:

Users should:

- Wash hands thoroughly before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Avoid spraying ponds or aquaria containing fish. Do not contaminate water by cleaning of equipment or when disposing of equipment wastewaters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency in your state responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, made of material such as barrier laminate or Viton
- Shoes plus socks

NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, or greenhouses. Keep people out of treated areas without protective clothing until sprays have dried.

GENERAL INFORMATION

RYCAR insecticide is formulated as a suspension concentrate containing 1.80 lbs. of active ingredient per gallon. **RYCAR** insecticide works primarily through contact action, and by ingestion, and is translaminar. This product is a contact insecticide used for the control of various insects. **RYCAR** insecticide should be used in a program with other products to provide season long protection. Apply as a spray as directed in the "Application Directions" section of this label.

RESISTANCE MANAGEMENT

RYCAR insecticide contains pyrifluquinazon, which has been classified by the Insecticide Resistance Action Committee (IRAC) as "unknown mode of action". Repeated use of the same chemistry has been shown to result in the buildup of resistant strains of insects. Rotate the use of **RYCAR** insecticide with alternate mode of action insecticides.

Resistance management strategies recommend that you DO NOT apply rates lower than recommended on the label. For best results, do not make more applications of **RYCAR** insecticide than permitted on the label in a single crop cycle. Contact your local extension specialist or certified crop advisor for additional resistance management or IPM recommendations. Also, for more information about Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org>.

MIXING DIRECTIONS

Shake well before using. Read and follow all label directions for each tankmix product prior to any tank mixing with **RYCAR** insecticide. This product can be

mixed with other registered pesticides. The most restrictive use directions and precautions must be followed. No labeled dose rate should be exceeded.

RYCAR Insecticide Alone: Begin with clean equipment. Fill spray tank with ¾ of the amount of water needed for the intended application and then turn on agitation. Pour recommended amount of product on the surface of water in the spray tank. Add the remaining water volume to the spray tank with agitation running. Keep agitation running during filling and spraying operations. If spraying must be stopped before emptying the sprayer, resume agitation before spraying the remainder of the load.

RYCAR Insecticide Tank Mixtures: Begin with clean equipment. Fill spray tank with ¾ of the amount of water needed for the intended application and turn on agitation. If using a buffering agent, add after filling the tank with ¾ amount of water. Add the recommended amount of tankmix products in the following order while maintaining agitation:

- 1) products in water-soluble packets
- 2) wettable powders
- 3) water-dispersible granulars and/or soluble powders
- 4) flowable liquids (including RYCAR insecticide)
- 5) emulsifiable concentrates
- 6) adjuvants and/or oils
- 7) remaining amount of water to achieve the desired level

RYCAR Insecticide Tank Mixtures with Products in Water-Soluble Packaging: Follow all use directions as listed above under **RYCAR Insecticide Alone** with the following exception: add the desired number of water-soluble bags to the tank (if allowed by their label instructions) at the same time the **RYCAR** insecticide is added.

Note: If using products in water soluble packaging, do not tank mix with products that contain boron, chromium, or other micronutrients.

RYCAR insecticide is physically and biologically compatible with many registered pesticides, fertilizers or micronutrients. Contact your SePRO turf and ornamental specialist for advice when considering mixing **RYCAR** insecticide with other pesticides, fertilizers or micronutrients. If you have no experience with the combination you are considering, you should conduct a test to determine physical compatibility. To determine physical compatibility, add the recommended proportions of each chemical with the same proportion of water, as will be present in the chemical supply tank, into a suitable container, mix thoroughly and allow to stand for five minutes. If the combination remains mixed, or can be readily re-mixed, the mixture is considered physically compatible.

APPLICATION DIRECTIONS

USE LIMITATIONS AND PRECAUTIONS

Indoor Greenhouse Use on Ornamental Crops

- Do not apply this product through any type of irrigation system.
- Do not apply this product as a smoke or aerosol.
- Do not harvest cut flowers for 48 hours after spraying.
- Keep compost in sheltered compost area.
- Do not apply this product to plants used for food or animal feed.
- Do not apply to fruit trees and vines that will bear harvestable fruit within 12 months.

Indoor Greenhouse Use on Ornamental Crops

RYCAR insecticide is recommended for use in indoor greenhouses on several ornamental plants. The use directions of this product are based on the results of product testing programs on a wide variety of ornamental plants. However, it is impossible to test this product on all species and cultivars. The phytotoxicity of **RYCAR** insecticide has been assessed on a wide variety of common ornamental plants with no phytotoxic effects. However, not all plant species and their varieties and cultivars have been tested with possible tankmix combinations, sequential pesticide treatments, and adjuvants and surfactants. Local conditions also can influence crop tolerance and may not match those under which testing has been conducted. Therefore, before using **RYCAR** insecticide, test the product on a sample of the crop to be treated to ensure that a phytotoxic response will not occur as a result of applications.

RYCAR insecticide works primarily through contact and translaminar activity, so good spray coverage is necessary for control of listed insects on the label. Applications should be made immediately after the spray solution is prepared. Under severe insect pressure, use the maximum rates and the shorter spray interval as specified on the label. Dense foliage or excessive growth will often prevent adequate coverage; adjust spray volumes accordingly. Mix with sufficient water and apply as a foliar spray to obtain uniform coverage. Treat plants when pests are immature or at a susceptible stage and populations are building, before crop damage occurs.

Applications may be made with high or low volume spray equipment that provides thorough coverage of the plant. Apply with properly calibrated spray equipment. A wetting agent or other spray adjuvant, approved for use on the crop, may be added

to spray solutions according to the manufacturer's use instructions to achieve optimum control.

USE LIMITATIONS AND PRECAUTIONS FOR INDOOR GREENHOUSE USE ON ORNAMENTAL CROPS

- Do not apply this product through any type of irrigation system.
- Do not apply this product as a smoke or aerosol.
- Do not harvest cut flowers for 48 hours after spraying.
- Keep compost in sheltered compost area.
- Do not apply this product to plants used for food or animal feed.
- Do not apply to fruit trees and vines that will bear harvestable fruit within 12 months.

CONTROLLING PESTICIDE RUNOFF IN GREENHOUSES

Do not allow pesticide spray solution to run off outside of the application area. Do not use in green-houses that discharge irrigation runoff to surface water drains. If applied in an area with a non-permeable floor and a floor drainage system, excess application spray or irrigation runoff must be collected and subjected to secondary wastewater treatment.

List of Plant Species Tested for Tolerance to RYCAR Insecticide¹

| Common Name | Scientific Name |
|----------------------|-----------------------------|
| Begonia | <i>Begonia</i> sp. |
| Balsam | <i>Impatiens</i> sp. |
| New Guinea Impatiens | Impatiens New Guinea Hybrid |
| Petunia | <i>Petunia</i> sp. |
| Marigolds | <i>Tagetes</i> sp. |
| Vervain | <i>Verbena</i> sp. |
| Zinnia | <i>Zinnia</i> sp. |

¹ Local conditions can influence crop tolerance and may not match those under which these species were tested. Before using **RYCAR** insecticide, test the product on a small sample of the crop to be treated.

APPLICATION RATE CHART FOR RYCAR INSECTICIDE

| GREENHOUSE ORNAMENTAL PLANTS | | |
|--|---------------------------------|---|
| Pest Controlled | Dilution Rate FL OZ/100 Gallons | Use Directions |
| Whiteflies | 1.6 to 3.2 fl oz | <ul style="list-style-type: none"> • Apply no more than 10 gallons of spray solution per 1,000 sq ft per application. • Do not apply more than 2 applications per crop cycle. • Do not apply more than 12.8 fl oz per crop cycle. • Allow at least 10 days between applications. • Use in sufficient volume to obtain uniform plant coverage. 100 gallons of finished spray solution will typically cover 20,000 sq ft of greenhouse. • Do not harvest cut flowers for 48 hours after spraying. |
| Aphids Leafhoppers Chilli Thrips | 2.4 to 3.2 fl oz | |
| Mealybugs | 6.4 fl oz | |

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Nonrefillable plastic containers

For liquid dilutables (Less than 5 gallons):

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining content into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

STORAGE AND DISPOSAL *(continued)*

Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For any dilutable pesticide (Greater than 5 gallons or 50 pounds):

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple Rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable plastic containers

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

TERMS AND CONDITIONS OF USE

If terms of the following *Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies* are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, to the extent consistent with applicable law, use by the buyer or any other user constitutes acceptance of the terms under *Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies*.

WARRANTY DISCLAIMER

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation or the seller. To the extent consistent with applicable law, all such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent consistent with applicable law, SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the *Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer* or this *Limitation of Remedies* in any manner.

Protected by Patent Number 645535

©Copyright 2014 SePRO Corporation

®Rycar is a registered trademark of Nichino



SePRO Corporation
11550 North Meridian Street, Suite 600
Carmel, IN 46032 U.S.A.